

<https://helda.helsinki.fi>

Thyroid carcinomas that occur in familial adenomatous polyposis patients recurrently harbor somatic variants in APC, BRAF, and KTM2D

Nieminen, Taina Tuulikki

2020-03-01

Nieminen , T T , Walker , C J , Olkinuora , A , Genutis , L K , O'Malley , M , Wakely Paul E , J , LaGuardia , L , Koskenvuo , L , Arola , J , Lepistö , A , Brock , P , Yilmaz , A S , Eisfeld , A-K , Church , J , Peltomäki , P & de la Chapelle , A 2020 , ' Thyroid carcinomas that occur in familial adenomatous polyposis patients recurrently harbor somatic variants in APC, BRAF, and KTM2D ' , Thyroid , vol. 30 , no. 3 , pp. 380-388 . <https://doi.org/10.1089/thy.2019.0561>

<http://hdl.handle.net/10138/327265>

<https://doi.org/10.1089/thy.2019.0561>

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Original Article

Thyroid carcinomas that occur in familial adenomatous polyposis patients

recurrently harbor somatic variants in *APC*, *BRAF*, and *KTM2D*

Taina T. Nieminen, PhD^{1,2,3}, Christopher J. Walker, PhD^{1,2}, Alisa Olkinuora, MSc³, Luke K. Genutis, BS², Margaret O'Malley, BS^{4,5}, Paul E. Wakely Jr. MD⁶, Lisa LaGuardia, RN, BSN^{4,5}, Laura Koskenvuo, MD, PhD⁷, Johanna Arola, MD, PhD³, Anna H. Lepistö, MD, PhD⁷, Pamela Brock, MS, LGC⁹, Ayse Selen Yilmaz, MS¹⁰, Ann-Kathrin Eisfeld, MD², James M. Church, MD^{4,5}, Päivi Peltomäki, MD, PhD³ and Albert de la Chapelle, MD, PhD²

1. These two first authors contributed equally

2. Department of Cancer Biology and Genetics, The Ohio State University Comprehensive

Cancer Center, The Ohio State University, Columbus, Ohio, 43210, USA

3. Department of Medical and Clinical Genetics, Biomedicum Helsinki, P.O.Box 63,

00014 University of Helsinki, Helsinki, Finland

25 4. Department of Colorectal Surgery, Cleveland Clinical, Lakewood, Ohio, 44107,

26

27 USA

28

29 5. Sanford R. Weiss MD Center for Hereditary Colorectal Neoplasia, Cleveland Clinic,

30

31 Lakewood, Ohio, 44107, USA

32

33 6. Department of Pathology, The Ohio State University Wexner Medical Center, The

34

35 Ohio State University, Columbus, Ohio, 43210, USA

36

37 7. Department of Gastrointestinal Surgery, Abdominal Center, P.O.Box 340, 00029

38

39 Helsinki University Hospital and University of Helsinki, Helsinki, Finland

40 8. Department of Pathology, University of Helsinki and HUSLAB, P.O.Box 400, 00029

41

42 Helsinki, Finland

43

44 9. Department of Internal Medicine, The Ohio State University Wexner Medical

45

46 Center, Division of Human Genetics, The Ohio State University, Columbus, Ohio,

47 43210, USA

48

10. Department of Biomedical Informatics, The Ohio State University, Columbus, Ohio,

43210, USA

Email addresses: taina.nieminen@osumc.edu, christopher.walker@osumc.edu,
alisa.olkinuora@helsinki.fi, luke.genutis@osumc.edu, OMALLEM@ccf.org,
paul.wakely@osumc.edu, LAGUARL@cff.org, laura.koskenvuo@hus.fi, johanna.arola@hus.fi,
anna.lepisto@hus.fi, pamela.brock@osumc.edu, AyseSelen.Yilmaz@osumc.edu, ann-kathrin.eisfeld@osumc.edu, CHURCHJ@ccf.org, paivi.peltomaki@helsinki.fi,
albert.delachapelle@osumc.edu

Running Title: FAP associated thyroid cancer

Key words: Familial adenomatous polyposis, *APC*, papillary thyroid cancer, cribriform-morular variant, whole genome sequencing

Abstract

Background: Familial adenomatous polyposis (FAP) is a condition typically caused by pathogenic germline mutations in the *APC* gene. In addition to colon polyps, individuals with FAP have a substantially increased risk of developing papillary thyroid carcinoma (PTC). Little is known about the events underlying this association, and the prevalence of somatic “second-hit” mutations in *APC* is controversial.

Methods: Whole genome sequencing was performed on paired thyroid tumor and normal DNA from 12 FAP patients who developed PTC. Somatic mutation profiles were compared with clinical characteristics and previously sequenced sporadic PTC cases. Germline variant profiling was performed to assess the prevalence of variants in genes previously shown to have a role in PTC predisposition.

Results: All 12 patients harbored germline mutations in *APC*, consistent with FAP.

Seven patients also had somatic mutations in *APC*, and seven patients harbored somatic mutations in *KMT2D*, which encodes a lysine methyl transferase. Mutation of these genes is extremely rare in sporadic PTCs. Notably, only two of the tumors harbored the somatic *BRAF* p.Val600Glu mutation, which is the most common driver mutation found in sporadic PTCs.

Six tumors displayed cribriform-morular variant of PTC (PTC-CMV) histology, and all six of these had somatic mutations in *APC*. Additionally, 9 FAP-PTC patients had rare germline variants in genes that were previously associated with thyroid carcinoma.

Conclusions: Our data indicate that FAP-associated PTCs typically have distinct mutations compared to sporadic PTCs. Roughly half of the thyroid cancers that arise in FAP patients have somatic “second-hits” in *APC*, which is associated with PTC-CMV

histology. Somatic *BRAF* p.Val600Glu variants also occur in some FAP patients, a novel finding. We speculate that in carriers of heterozygous pathogenic mutations of tumor suppressor genes such as *APC*, a cooperating second-hit somatic variant may occur in a different gene such as *KTM2D* or *BRAF*, leading to differences in phenotypes. The role of germline variance in genes other than *APC* (9 of 12 patients in this series) needs further research.

Introduction

In addition to colon polyps and colorectal cancer, patients with familial adenomatous polyposis (FAP) have an increased risk to develop extracolonic malignancies and benign conditions/tumors. Of particular interest is the frequent occurrence of papillary thyroid cancer (PTC) which is some 100 times more prevalent in FAP patients than in the general population (1). FAP is typically caused by germline mutations in the *APC* gene that result in a truncated APC protein, and inevitably develops into colorectal cancer (CRC) when somatic “second hit” mutations in *APC* occur in colon cells (2). Previous studies have confirmed the existence of such “second hits” in the *APC* gene in a subset of thyroid cancers that occur in FAP patients, (3) but this remains controversial, as other reports have concluded that the *APC* gene is rarely somatically mutated in FAP-associated PTC (4,5,6). This contradiction has not been resolved.

FAP-associated PTCs often display cribriform-morular variant (CMV) histology, which is otherwise extremely rare (~0.2% of all thyroid cancers) (7). Altogether about half of all CMV-PTCs occur in FAP patients (7,8). In general PTC is about three times more common in females than in males, and this ratio is even higher in PTC-CMV patients and FAP-associated PTCs (7). Molecular characteristics of PTC-CMV include mutations in the *CTNNB1* and/or *PIK3CA* genes and *RET/PTC* rearrangements. No oncogenic *BRAF* mutations have so far been reported in CMV-PTCs or FAP-associated PTCs; however a comprehensive assessment of the somatic alterations that occur in FAP-associated PTCs has not been conducted. To better understand the germline and somatic variants found in this unique tumor type and to try to learn more about the genetic mechanisms of thyroid cancer development in individuals with FAP-associated germline mutations in *APC*, we performed whole genome sequencing of paired tumor

and normal DNA from 12 FAP-associated PTC patients.

Material and methods

Patients

All patients with both FAP and PTC were selected from the PTC patient repositories at

the Cleveland Clinic and the Helsinki University Hospital. Histological review of tumor

sections was performed to confirm the presence of cancer in the resected thyroid. This

study was limited to patients with germline *APC* mutations detected by clinically-

approved testing methods. All patients provided written informed consent and studies

were performed in accordance with the declaration of Helsinki, and approved by institutional

review boards at both institutions.

DNA extraction and sequencing

DNA was extracted from paraffin embedded thyroid tumors and adjacent normal tissue

(patients 2, 4, 5, 7, 8, 9, 10, and 12) using QIAamp DNA FFPE Tissue Kits (Qiagen,

Hilden, Germany), or from blood (patients 1, 3, 6, and 11) using a previously described

non-enzymatic DNA-extraction method (9). DNA samples were quantified using a Qubit

3.0 Fluorometer (Thermo Fisher Scientific, Waltham, MA) and fragment size was

assessed using a 2100 Bioanalyzer system (Agilent Technologies, Santa Clara, CA). All

samples had average fragment sizes > 1500bp. Library preparation and paired-end

genome sequencing was performed by Novogene (Beijing, China), using Truseq Nano DNA HT

Sample Preparation Kits (Illumina, San Diego, CA) and HiSeq 4000 instruments (Illumina, San

Deigo, CA). All samples were sequenced to > 80 gigabytes of data.

Informatics

The quality of sequences was confirmed using FASTQC software. All samples had >99% of reads mapped, and all samples had >90% of bases with phred-scaled quality scores > 30. Mapping was done with Burrows-Wheeler Aligner to the Genome Reference Consortium Human Build 37. Samples were sorted with SamTools, duplicates were marked with Picard, and variants were called with GATK then annotated with ANNOVAR. For somatic and germline variant analysis, paired .vcf files from each tumor and matching adjacent normal thyroid or blood sample were loaded into BasePlayer software (10). Variants present in the tumor sample and absent but covered in the paired germline DNA were considered somatic. Publicly available data from the American Association for Cancer Research Genomics Evidence Neoplasia Information Exchange (AACR-GENIE) and The Cancer Genome Research Atlas (TCGA) were accessed using cbiportal.org on August 19, 2019.

Somatic variant signature analysis was performed using the R package

DeconstructSigs,(11) and the 30 signatures described in the Catalog of Somatic

Mutations in Cancer (COSMIC) Mutational Signatures (v2-March 2015).

Somatic loss of heterozygosity (LOH) was determined using BasePlayer software (10)

by comparing the allelic ratios of sequenced germline variants to their corresponding

ratios in the matched tumor sample as described (12,13). A tumor to normal ratio of

≤ 0.6 or ≥ 1.67 was considered LOH, and a tumor to normal ratio of 0.6-0.8 or 1.25-1.67

was considered putative LOH

Results

The 12 FAP-PTC patients in our cohort consisted of 4 males and 8 females (Table 1).

Five patients displayed classic PTC histology (three males and two females), six

patients had PTC-CMV (one male and five females), and one female had follicular

variant of papillary thyroid carcinoma. Number of the tumors varies from 1 to 2 per individual

and tumor sizes varies between 0.1 cm and ? All patients in our sample set were diagnosed

with FAP before thyroid cancer, with an average age at FAP diagnosis of 25 years

(range 12-44 years), and an average age at PTC diagnosis of 38 years (range 20-62

years). Patients were diagnosed with thyroid carcinoma on average 15 years after

being diagnosed with FAP (range 1-46 years after FAP diagnosis). This cohort

displayed characteristics typically associated with FAP, including colorectal cancer,

ampullary cancer, and desmoid tumors.

Whole genome sequencing was performed on paired tumor and normal DNA from all

12 cases. An oncoprint of the most commonly mutated genes is presented in Figure 1a.

Seven patients had somatic “second-hit” mutations in *APC*, including one patient with

clear somatic LOH of *APC* and two patients with putative LOH. The lysine methyl

transferase gene *KMT2D* was also mutated in seven of the 12 PTC tumors, and

another *KMT2* family member, *KMT2C* was mutated in three patients (four mutations). The

recurrently mutated genes in these samples were strikingly different from genes that are

typically mutated in sporadic PTC, as evidenced by comparison with the AACR-GENIE and

TCGA of PTC data sets (Supplementary Table 1). Specifically, mutations in *KMT2C*,

KMT2D and *APC* were only detected in <3% of sporadic cases. Conversely, *BRAF* is

mutated in about 60% of all PTCs, and the vast majority of these mutations consist of the

known cancer driver mutation *BRAF* p.Val600Glu. However, in our sample set only two

cases harbored *BRAF* mutations (both p.Val600Glu). Notably, neither of these samples

had somatic *APC* mutations. *RET*/PTC translocations are also common in typical

PTCs,(14) however, in our cohort, there was no evidence of breakpoints between exons

11 and 12 in *RET*. Therefore, it is likely that none of the APC-associated PTCs we

sequenced harbored *RET*/PTC or other *RET* translocations/inversions. We did not detect

any somatic mutations in the Ras genes (*NRAS*, *HRAS* and *KRAS*), which are recurrently

mutated in sporadic PTC.

All clinically detected germline nonsense, frameshift and insertion/deletion variants in

the *APC* gene were validated in our genome sequencing. Three samples had complex

APC germline mutations: patient 11 had a heterozygous ~23Mb deletion of

chromosome 5q that contained *APC*, patient 4 had a deletion of exons 15-16

(NM_000038, coding exons 14-15), and patient 1 had a C to T transition within intron 11

that results in incorporation of an out of frame pseudoexon and underlies FAP, as we

have previously described (Table 2) (15). The remaining nine patients all had frameshift and nonsense germline mutations in the *APC* coding sequence, located between codons 471 and 1465 (Fig. 1b, Table 2). The “second-hit” somatic variants in *APC* were more spread out, and there was not an obvious correlation between the location of the germline mutation in *APC* and the somatic second hit, in contrast to reports that the location of the germline FAP-associated *APC* mutation can influence the position of the somatic *APC* variant in colorectal cancer secondary to FAP (Fig. 1b, Table 2) (16).

We examined the specific base pair substitutions in the mutations in these samples and compared them to described mutational signatures associated with different cancers and cancer subtypes (17). Most samples showed strong correlations with expression signatures 1 (pan-cancer deamination of 5-methylcytosine), 3 (failure of DNA double-

strand break-repair by homologous recombination), 5 (pan-cancer with unknown
etiology), 12 (unknown etiology signature found in liver cancer) and 20 (defective DNA
mismatch repair) (Fig. 2a, b). Patient 6's tumor exhibited a strong signal for mutation
signature 6, which is seen in microsatellite unstable tumors that have DNA mismatch
repair defects, and mutation signature 19, which is an unknown etiology signature found
in pilocytic astrocytomas (Fig. 2c) (17,18,19,20). Notably, patient 6 did not have
somatic mutations in *MLH1*, *PMS2*, *MSH2*, or *MSH6*, and also did not have likely
pathogenic germline variants in these genes.

Finally we examined the germline variants in these samples for alleles that might play a
role in thyroid cancer, specifically focusing on nonsynonymous variants in genes
previously implicated in familial thyroid cancer (21). Because rare
variants are more likely to be high-risk alleles for diseases than common variants, based

on family studies (22), we only examined nonsynonymous variants with a population minor allele frequency less than 0.01 in the gnomAD database. We found 9 of the FAP-PTC patients had germline variants in 17 different genes, which were previously found to harbor variants associated with familial thyroid carcinoma (21) (Supplementary Table 2). Interestingly, there were several variants in the *RNF213* gene in three different FAP-PTC patients. *RNF213* has been found to be mutated in liver cancer (23) and all three of the patients (4, 5, 8) with *RNF213* variants had a strong signal for mutational signature 12 (Fig. 2a), which is linked to liver cancer. The p.Arg752Leu *FGD6* variant was found in two patients, one from the USA (patient 9) and one from Finland (patient 1). *FGD6* is located on chromosome 12q22, and this band has been observed to be amplified in thyroid adenomas, and might also play a role in thyroid carcinomas as well (24).

Discussion

It is surprising, although not entirely unexpected, that we detected somatic second hits in *APC* in over half of the FAP-PTC tumors we analyzed. Cetta et al. reported that somatic mutations in *APC* do not occur in FAP-associated PTC, (4,25) but in contrast, somatic second hit mutations in *APC* were previously reported by other groups (3,5,6).

One explanation of this apparent contradiction is the recently improved sequencing methodology, and the failure of some researchers to examine the entire *APC* coding sequence.

Mechanistically, pathogenic nonsense and frameshift *APC* mutations lead to truncated *APC* protein products that are unable to interact with the cytoplasmic complex that mediates β -catenin degradation. Thus, the β -catenin/Lef/Tcf complex remains unchecked in the nucleus where it activates WNT signaling pathways responsible for

enhanced cellular migration, proliferation and loss of differentiation (26). Our finding is

consistent with the idea that somatic second hits and/or LOH in *APC* further add

cancerous properties to the cell and likely contribute to malignant transformation.

Our identification of the somatic *BRAF* p.Val600Glu mutation in two patients is a novel

finding, as to our knowledge, no single case has been described in the literature where

oncogenic *BRAF* mutations occur in either FAP-PTC or PTC-CMV

(7,27,28,29,30,31,32). This implies that some PTCs arising in the context of germline

pathogenic *APC* variants can share the same driver mutations as sporadic PTCs. The

mutual exclusivity of the somatic *APC* and *BRAF* mutations is consistent with different

molecular subtypes of PTC occurring in different FAP patients. The tumors with *BRAF*

p.Val600Glu mutations displayed typical PTC histology, and occurred in one male and

one female. However, the patients with non-silent somatic *APC* variants were almost

entirely female (6 to 1, female to male), and in six of seven cases showed PTC-CMV

histology. Interestingly, the patient with a somatic *APC* mutation who did not have PTC-CMV

histology (patient 7) harbored the most 3' *APC* mutation we detected. The mutated *APC*

protein in patient 7 likely retains some beta-catenin binding ability, and we speculate this

could contribute to why patient 7 did not have CMV histology.

What does it mean that 58% and 33% of the PTC tumors in these FAP patients have

somatic variants in the *KMT2D* and *KMT2C* genes, respectively? *KMT2D* and *KMT2C*

are methyl transferase genes that encode important pieces of the COMPASS complex

(33). Pathogenic somatic mutations in both of these genes have been detected in many

different cancers, such as oropharyngeal squamous cell carcinoma, T-cell lymphoma,

bladder cancer, head and neck cancer, and breast and endometrial cancers, but are

extremely rare in sporadic PTC (34,35,36,37). Notably, *KMT2D* somatic variants have

498
499 been shown to contribute to increased mutational burden and genome instability (38).
500

501 Prompted by *KMT2D* somatic variants, we looked for biological link between *APC* and *KMT2D*
502

503 genes, and to our surprise, it seems that also *KMT2D* might be involved in WNT signaling.
504

505 *KMT2D* together with *ALK* gene are connected with *CTNNB1* (β -catenin) (Pinckney et al. 2018,
506

507 Applied Cancer Research, 38:13), so it is not surprise that in some tumors instead of *APC*,
508

509 somatic mutations occurs in *KMT2D*.
510

511 We speculate that the detected somatic variants in these genes
512

513 (particularly *KMT2D*) might be important in the context of deactivated WNT signaling caused by
514

515 FAP-associated germline *APC* mutations, and evidence of epigenetic dysregulation in these
516

517 cases warrants further investigation. Our data implicate that the mutations in *KMT2D* may be
518

519 cancer drivers in the patients in which they were observed.

The overall female to male ratio in our cohort is less skewed towards

females than other reports of FAP-PTC patient demographics (7,39). For example, Lam et

al.(7) reported that the female to male ratio in PTC-CMV is 31 to 1, whereas in our cohort

there was only a 5 to 1 ratio of females to males among patients with PTC-CMV.

However, we do acknowledge that the modest size of our cohort does not lend itself to

definitive conclusions regarding sex ratios. Seven of the twelve patients had desmoid

tumors, which is not surprising given that this a common feature of FAP patients

(29,40,41,42,43,44).

Our study is the first that suggests mutations in genes other than *APC* can cooperate with the

germline *APC* variant in FAP patients to drive thyroid cancer. Also, our

findings provide new information on the genetic steps that participate in the carcinogenesis

process. Our data are consistent with a model where the pathogenic germline *APC* variants act as “gatekeepers” in the thyroid. Some patients, almost always females, acquire somatic second hits in *APC* that drive a thyroid cancer with CMV histology. In other cases, oncogenic activating mutations somatically occur in *BRAF*, similar to sporadic thyroid cancers. In patients who lack clear driver mutations in *APC* or *BRAF*, an intriguing possibility is that somatic variants in other genes (e.g. *KTM2D*, *KMT2C* and others) may act as cancer drivers in the thyroid. This concept postulates that a somatic heterozygous variant in a gene such as *KTM2D* can act as a trigger of the malignant transformation of a cell heterozygous for pathogenic variant in another gene (i.e. *APC*), and is in line with the concept that multiple events contributing different cancerous properties to a cell need to occur in order for a malignancy to develop and proliferate (45). To prove or disprove this scenario in FAP-associated thyroid cancer, a larger series of

cases will need to be studied, and gene functions and interactions carefully

documented.

It is striking that 9 out of the 12 patients we sequenced harbored rare nonsynonymous

mutations in 17 selected genes known to be associated with familial thyroid cancer. This

is consistent with the idea that additional germline variants other than of *APC* can contribute

to PTC formation in FAP patients. Further studies are necessary to unequivocally prove

a causative role for the implicated germline variants in FAP-associated PTC, and explore

their interactions with the altered WNT signaling caused by pathogenic *APC* variants.

Acknowledgements

The authors would like to acknowledge: the patients who consented to provide material

for this study; Jan Lockman and Barbara Fersch for administrative help; the OSUCCC

Genomics Shared Resource and the Plant Microbe Genomics Facility for sequencing assistance; the OSUCCC Human Cancer Genetics Sample Bank for sample processing and storage; and the Ohio Supercomputer Center for computational resources. This work was supported by the Leukemia Clinical Research Foundation, the Pelotonia fellowship program, the Jane and Aatos Erkkö Foundation, the Finnish Cancer Organizations, the Sigrid Juselius Foundation, the Inkeri and Mauri Vänskä Foundation, the HiLIFE Fellows 2017-2020, the National Cancer Institute Grants P01CA124570 and P50CA168505, and The Academy of Finland Grant294643.

Author Disclosure Statement:

The authors declare no conflict of interest.

Correspondence: Taina Nieminen, 850 Biomedical Research Tower, 460 W 12th Ave, Columbus, OH 43210, email: taina.nieminen@osumc.edu, ph: (614)-843-8913

References

1. Herraiz M, Barbesino G, Faquin W, Chan-Smutko G, Patel D, Shannon KM, Daniels GH, Chung DC 2007 Prevalence of thyroid cancer in familial adenomatous polyposis syndrome and the role of screening ultrasound examinations. Clin Gastroenterol Hepatol 5:367-373.
2. Nallamilli BR, Hegde M 2017 Detecting *APC* gene mutations in familial adenomatous polyposis (FAP). Curr Protoc Hum Genet 92:10.18.11-10.18.16.
3. Soravia C, Sugg SL, Berk T, Mitri A, Cheng H, Gallinger S, Cohen Z, Asa SL, Bapat BV 1999 Familial adenomatous polyposis-associated thyroid cancer: a clinical, pathological, and molecular genetics study. Am J Pathol 154:127-135.
4. Cetta F, Dharmo A, Malagnino G, Barellini L 2007 Germ-line and somatic mutations of the *APC* gene and/or β -catenin gene in the occurrence of FAP associated thyroid carcinoma. World J Surg 31:1366-1369.
5. Iwama T, Konishi M, Iijima T, Yoshinaga K, Tominaga T, Koike M, Miyaki M 1999 Somatic mutation of the *APC* gene in thyroid carcinoma associated with familial adenomatous polyposis. Jpn J Cancer Res 90:372-376.
6. Uchino S, Noguchi S, Yamashita H, Yamashita H, Watanabe S, Ogawa T, Tsuno A, Murukami A, Miyauchi A 2006 Mutational analysis of the *APC* gene in cribriform-morular variant of papillary thyroid carcinoma. World J Surg 30:775- 779.
7. Lam AK, Saremi N 2017 Cribriform-morular variant of papillary thyroid carcinoma: a distinctive type of thyroid cancer. Endocr Relat Cancer 24:R109- R121.

8. Uchino S, Ishikawa H, Miyauchi A, Hirokawa M, Noguchi S, Ushima M, Yoshida T, Michikura M, Sugano K, Sakai T 2016 Age- and gender-specific risk of thyroid cancer in patients with familial adenomatous polyposis. *J Clin Endocrinol Metab* 101:4611-4617.
9. Lahiri DK, Nurnberger JI Jr 1991 A rapid non-enzymatic method for the preparation of HMW DNA from blood for RFLP studies. *J Nucl Acids Res* 19:5444.
10. Katainen R, Donner I, Cajuso T, Kaasinen E, Palin K, Makinen V, Aaltonen LA, Pitkanen E 2018 Discovery of potential causative mutations in human coding and noncoding genome with the interactive software BasePlayer. *Nat Protoc* 13:2580-2600.
11. Rosenthal R, McGranahan N, Herrero J, Taylor BS, Swanton C 2016 DeconstructSigs: delineating mutational processes in single tumors distinguishes DNA repair deficiencies and patterns of carcinoma evolution. *Genome Biol* 17:31.
12. Porkka N, Valo S, Nieminen T, Olkinuora A, Maki-Nevala S, Eldfors S, Peltomaki P 2017 Sequencing of Lynch syndrome tumors reveals the importance of epigenetic alterations. *Oncotarget*. 14:108020-108030.
13. Ollikainen M, Abdel-Rahman WM, Moisio AL, Lindroos A, Kariola R, Jarvela I, Poyhonen M, Butzow R, Peltomaki P 2005 Molecular analysis of familial endometrial carcinoma: a manifestation of hereditary nonpolyposis colorectal cancer of a separate syndrome? *J Clin Oncol* 20:4609-4616.
14. Cetta F, Chiappetta G, Melillo RM, Petracci M, Montalto G, Santoro M, Fusco A 1998 The ret/ptc1 oncogene is activated in familial adenomatous polyposis- associated thyroid papillary carcinomas. *J Clin Endocrinol Metab* 83:1003-1006.
15. Nieminen TT, Pavicic W, Porkka N, Kankainen M, Jarvinen HJ, Lepisto A, Peltomaki

- 666 P 2016 Pseudoxons provide a mechanism for allele-specific expression of APC in
667 familial adenomatous polyposis. *Oncotarget* 7:70685- 70698.
- 668 16. Albuquerque C, Breukel C, van der Luijt R, Fidalgo P, Lage P, Slors FJ, Leitao CN,
669 Fodde R, Smits R 2002 The 'just-right' signaling model: APC somatic mutations are
670 selected based on a specific level of activation of the beta-catenin signaling cascade.
671 *Hum Mol Genet* 11:1549-1560.
- 672 17. Alexandrov LB, Nik-Zainal S, Wedge DC, Campbell PJ, Stratton MR 2013 Deciphering
673 signature of mutational processes operative in human cancer. *Cell Rep* 3:246-259.
- 674 18. Nik-Zainal S, Alexandrov LB, Wedge DC, Van Loo P, Greenman CD, Raine K, Jones
675 D, Hinton J, Marshall J, Stebbings LA, Menzies A, Martin S, Leung K, Chen L, Leroy C,
676 Ramakrishna M, Rance R, Lau KW, Mudie LJ, Varela I, McBride DJ, Bignell GR,
677 Cooke SL, Shlien A, Gamble J, Whitmore I, Maddison M, Tarpey PS, Davies HR,
678 Papaemmanuil E, Stephens PJ, McLaren S, Butler AP, Teague JW, Jönsson G,
679 Garber JE, Silver D, Miron P, Fatima A, Boyault S, Langerød A, Tutt A, Martens JW,
680 Aparicio SA, Borg Å, Salomon AV, Thomas G, Børresen-Dale AL, Richardson AL,
681 Neuberger MS, Futreal PA, Campbell PJ, Stratton MR; Breast Cancer Working
682 Group of the International Cancer Genome Consortium 2012 Mutational processes
683 molding the genomes of 21 breast cancers. *Cell* 151:979-993.
- 684 19. Helleday T, Eshtad S, Nik-Zainal S 2014 Mechanisms underlying mutational
685 signatures in human cancers. *Nat Rev Genet* 15:585-598.
- 686 20. Alexandrov LB, Stratton MR 2014 Mutational signatures: the patterns of somatic
687 mutations hidden in cancer genomes. *Curr Opin Genet Dev* 24:52-60.
- 688 21. Wang Y, Liyanarachchi S, Miller KE, Nieminen TT, Comiskey DF, Li W, Brock P,
689 Symer DE, Akagi K, He H, Koboldt DC, de la Chapelle A 2019 Whole genome

sequencing reveals rare germline variants leading to familial non-medullary thyroid cancer. Apr. 8.

22. Foulkes WD 2008 Inherited susceptibility to common cancers. *N Eng J Med* 359:2143-53.

23. Li X, Xu W, Kang W, Wong SH, Wang M, Zhou Y, Fang X, Zhang X, Yang H, Wong CH, To KF, Chan SL, Chan MTV, Sung JJY, Wu WKK, Yu J 2018 Genomic analysis of liver cancer unveils novel driver genes and distinct prognostic features. *Theranostics* 12;8:1740-1751.

24. Liu Y, Cope L, Sun W, Wang Y, Prasad N, Sangenario L, Talbot K, Somervell H, Westra W, Bishop J, Califano J, Zeiger M, Umbricht C 2013 DNA Copy number variations characterize benign and malignant thyroid tumors. 98:E558-66.

25. Cetta F, Pelizzo MR, Curia MC, Barbarisi A 1999 Genetics and clinicopathological findings in thyroid carcinomas associated with familial adenomatous polyposis. *Am J Pathol* 155:7-9.

26. Gao C, Wang Y, Broaddus R, Sun L, Xue F, Zhang W 2018 Exon 3 mutations of *CTNNB1* drive tumorigenesis: a review. *Oncotarget* 9:5492-5508.

27. Rossi ED, Revelli L, Martini M, Taddei A, Pintus C, Panunzi C, Fadda G 2012 Cribriform-morular variant of papillary thyroid carcinoma in an 8-year-old girl: a case report with immunohistochemical and molecular testing. *Int J Surg Pathol* 20:629-632.

28. Nakazawa T, Celestino R, Machado JC, Cameselle-Teijeiro JM, Vinagre J, Eloy C, Benserai F, Lameche S, Soares P, Sobrinho-Simoes M 2013 Cribriform-morular variant of papillary thyroid carcinoma displaying poorly differentiated features. *Int J Surg Pathol* 21:379-389.

29. Giannelli SM, McPhaul L, Nakamoto J, Gianoukakis AG 2014 Familial adenomatous

polyposis-associated, cribriform morular variant of papillary thyroid carcinoma harboring a K-RAS mutation: case presentation and review of molecular mechanisms. *Thyroid* 24:1184-1189.

30. Kwon MJ, Rho YS, Jeong JC, Shin HS, Lee JS, Cho SJ, Nam ES 2015 Cribriform-morular variant of papillary thyroid carcinoma: a study of 3 cases featuring the *PIK3CA* mutation. *Hum Pathol* 46:1180-1188.

31. Brehar AC, Terzea DC, Ioachim DL, Procopiu C, Brehar FM, Bulgar AC, Ghemigian MV, Dumitrache C 2016 Cribriform-morular variant of papillary thyroid carcinoma at pediatric age - case report and review of the literature. *Rom J Morphol Embryol* 57:531-537.

32. Oh EJ, Lee S, Bae JS, Kim Y, Jeon S, Jung CK 2017 TERT promoter mutation in an aggressive cribriform morular variant of papillary thyroid carcinoma. *Endocrine Pathology* 28:49-53.

33. Lang, A, Yilmaz M, Hader C, Murday S, Kunz X, Wagner N, Wiek C, Petzsch P, Köhrer K, Koch J, Hoffmann MJ, Greife A, Schulz WA 2019 Contingencies of UTX/KDM6A action in urothelial carcinoma. *Cancers (Basel)* 11:481.

34. Haft S, Ren S, Xu G, Mark A, Fisch K, Guo TW, Khan Z, Pang J, Ando M, Liu C, Sakai A, Fukusumi T, Califano JA 2019 Mutation of chromatin regulators and focal hotspot alterations characterize human papillomavirus-positive oropharyngeal squamous cell carcinoma. *Cancer* 125:2423-2434.

35. Fernandez-Pol S, Ma L, Joshi RP, Arber DA 2019 A survey of somatic mutations in 41 genes in a cohort of T-cell lymphomas identifies frequent mutations in genes involved in epigenetic modification. *Appl Immunohistochem Mol Morphol* 27:416-422.

36. Herz HM. Enhancer deregulation in cancer and other diseases 2016 *Bioessays*

38:1003-1015.

37. Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, Sun Y, Jacobsen A, Sinha R, Larsson E, Cerami E, Sander C, Schultz N 2013 Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. *Sci Signal* 6:pl1.
38. Kantidakis T, Saponaro M, Mitter R, Horswell S, Kranz A, Boeing S, Aygün O, Kelly GP, Matthews N, Stewart A, Stewart AF, Svejstrup JQ 2016 Mutation of cancer driver *MLL2* results in transcription stress and genome instability. *Genes Dev* 15:408-420.
39. Cetta F 2015 FAP associated papillary thyroid carcinoma: a peculiar subtype of familial nonmedullary thyroid cancer. *Patholog Res Int* :309348.
40. Chong J, Koshiishi N, Kurihara K, Kubuno S, Kawai T, Fukayama M 2000 Aspiration and imprint cytopathology of thyroid carcinoma associated with familial adenomatous polyposis. *Diagn Cytopathol* 23:101-105.
41. Dalal KM, Moraitis D, Iwamoto C, Shaha AR, Patel SG, Ghossein RA 2006 Clinical curiosity: cribriform-morular variant of papillary thyroid carcinoma. *Head Neck* 28:471-476.
42. Crippa S, Saletti P, Barizzi J, Mazzucchelli L 2012 The clinical management in familial adenomatous polyposis deserves continuous monitoring for thyroid carcinoma. *BMJ Case Rep*:bcr2012007046.
43. Abdullah Suhaimi SN, Nazri N, Nani Harlina ML, Md Isa N, Muhammad R 2015 Familial adenomatous polyposis-associated papillary thyroid cancer. *Malays J Med Sci* 22:69-72.
44. Alikhan M, Koshy A, Hyjek E, Stenson K, Cohen RN, Yeo KT 2015 Discrepant serum and urine β -hCG results due to production of β -hCG by a cribriform- morular variant of thyroid papillary carcinoma. *Clin Chim Acta* 438:181-185.

762 45. Hanahan D, Weinberg RA 2011 Hallmarks of cancer: The next generation. Cell 4:646-

763 674.

764

765

Table 1. Clinical characteristics of the 12 PTC patients with FAP

Patient no.	Sex	Histology	Age (FAP)	Age (thyroid cancer)	Number thyroid tumors	Tumor size(s)	Ascertainment	Other FAP-related disease presentation
1	Male	PTC-CMV	12	30	1	9 cm	Incidental	desmoid tumors
2	Female	PTC-CMV	21	35	2	0.2 cm 0.1 cm	Screening ultrasound	desmoid tumors
3	Female	PTC-CMV	43	45	1	2.4 cm	Mass in the neck, "self finding"	.
4	Female	PTC-CMV	33	35	2	0.9 cm 0.3 cm	Screening ultrasound	colon cancer
5	Female	PTC-CMV	18	43	2	0.9 cm 0.2 cm	Screening ultrasound	.
6	Female	PTC-CMV	21	21	2	2 cm, 3 cm	Difficulties to swallow	desmoid tumors
7	Female	PTC	18	42	1	0.9 cm	Screening ultrasound	unspecified extrathyroid cancer
8	Male	PTC	20	44	2	0.8 cm and smaller	Screening ultrasound	desmoid tumors, adrenal adenoma
9	Female	PTC	16	62	1	0.4 cm	Screening ultrasound	ampullary cancer
10	Male	PTC	23	28	1	1.6 cm	Screening ultrasound	desmoid tumors
11	Female	PTC-FV	24	24	1	0.9 cm	Jugular vein thrombus	desmoid tumors
12	Male	PTC	44	45	2	0.6 cm 0.4 cm	Screening ultrasound	desmoid tumors

Additional file 4: **Figure S1.** ALK and KMT2D (MLL2) converge on CTNNB1. An evaluation of the potential role of the gene products from mutated genes in our cohort that are associated with negative outcomes was completed to potentially identify pathways that could be targeted for novel therapy in this young group. String®, a protein-protein network modeling website sponsored by the String Consortium and the Swiss Institute of Bioinformatics (SIB) [], was used to postulate how the gene mutations were connected in biochemical pathways that could potentially create the carcinogenic phenotype. In our cohort, ALK and KMT2D (MLL2) gene mutations were found to be associated with both ovarian and endometrial negative outcomes in patients with neuroendocrine histology. In the String program, the ALK and KMT2D proteins, while being involved in multiple pathways, appear to converge on CTNNB1, a key member of the Wnt pathway. The Wnt pathway is important in cell adhesion and maintenance of an appropriate cell cycle and has been found to be compromised in a variety of cancers. Additionally, dysregulation of the Wnt pathway has recently been implicated in maintenance of cancer stem cells, metastasis and immune control (Zhan T et al., Oncogene 2017; 36:1461–1473). The String algorithm also revealed that HSP90AA1 is also involved in this complex pathway. HSP90AA1 has been shown to be a prognostic indicator of both liver and breast cancers. () (DOCX 687 kb)